## THIS IS AN URGENT CALL FOR RESEARCHERS AND DOCTORS IN HOSPITALS AND UNIVERSITIES TO INCREASE TESTING OF THE IMMUNE SYSTEM AFTER FIRST AND SECOND JABS

Patients were tested before and after Covid 19 vaccination on 3 separate occasions. Blood tests showed that after the Covid 19 vaccine, the white blood cell levels in all three patients suffered a major decline.

## After the first vaccination

CD8 killer T cells show a massive increase, as they kill cells infected with the spike protein

## After the second vaccination

The % of granulocytes jumps to 79.6%. This is mainly comprised of neutrophils, whose job is to clean up tissue damage. But what is being damaged inside the body so much?

Lymphocyte numbers fall to 13.8% - the lowest end of the normal range.CD4, CD8, B-lymphocytes and natural killer cells are all massively diminished.

	Before	After
Granulocyte %	58.9%	79.6%
Lymphocyte %	33.8%	13.8%

"I would look at this lab, and in 2 seconds I would say This person has auto-immunity. Let me say that again. This person has auto-immunity. People with auto-immunity – so many times I have seen clinically – are people with high granulocyte levels and a tanked adaptive immune system; the very thing designed to handle cancer cells and virally infected cells has absolutely tanked.

How on earth can they ever say this is working?

If this persists, their adaptive immune system will be very susceptible to any other viral infection – leaving them wide-open to 'break-through' infections"

If you are a doctor, please, please start testing their immune systems after the first and second jabs.

The test used is very similar to how we measure the progression of the HIV infection – measuring CD4, CD8, B-lymphocyte and Natural Killer T cell levels.